

Purification of water and air is promoting global warming and country decline

Shoichiro Ozaki

The Institute of Physical and Chemical Research 2-1 Hirosawa, Wakoshi Saitama Japan

Corresponding author Tel and Fax +81 0467670991

E-mail ozaki-0991@jcom.zaq.ne.jp

Abstract



Burning of fossil is increasing. Production of CO₂ and NO_x is increasing. Increased CO₂ and NO_x promoted the CO₂ assimilation. Most produced CO₂ is fixed by CO₂ assimilation. But developed countries started purification of water and air by elimination of NO_x and NP at around 1980. 6 billion tone NO_x and 2 billion tone NP are eliminated. NO_x is main nitrogen fertilizer and NP is main nitrogen and phosphorous fertilizer. Therefore plant growth is retarded. CO₂ fix is retarded. CO₂ is increasing. Food like grain, fish ,meat production is retarded. DGP increase rate decreased. Global warming and country decline are progressing. If developed countries stop NO_x elimination by ammonia and close waste water purification station, global warming will stop and country decline will stop.

Keyword:- purification of water, purification of air , GWPR, global warming protection ratio , plankton, NO_x elimination, NP elimination,CO₂ assimilation

Introduction

The earth is warmed by the fossil fuel burning releasing CO₂ and heat. The plant is growing by CO₂ assimilation absorbing CO₂ and heat producing carbohydrate and oxygen.

CO₂ assimilation



Burning

GWPR(Global warming protection ratio) = Produced CO₂/Fixed CO₂

If we can compensate the generation of CO₂ and heat with the generation of CO₂ and heat with the absorption of CO₂ and heat by CO₂ assimilation, GWPR(global warming protection ratio) become 1, and global warming can be protected.

About 510 billion tone CO₂ is produced by burning of fossil and respiration of animals.. About 30% of produced CO₂ is fixed by land plant CO₂ assimilation at land. About 70 % of produced CO₂ is fixed by plankton CO₂ assimilation at sea.

CO₂ concentration is increasing 2ppm every year. 140 billion tone CO₂ is increasing every year. Fixed CO₂ is 370 billion tone. Therefore global warming protection ratio is 510/ 370 = 1.38 We must decrease produced CO₂ and increase fixed CO₂ to lower GWPR. To increase fix of CO₂, we must increase CO₂ assimilation. To increase CO₂ assimilation, we must increase the supply of NP. we must increase NP concentration of sea .

CO₂ assimilation by plankton is most important reaction to control climate.. Plankton grow by eating CO₂,H₂O, nitrogen and phosphorous by Redfield ratio C: N: P 105.4 : 16 : 1 or 6.6 : 1 : 0.06. Plankton ask more N and P than normal plant. Ratio C: N: P 25 : 1: 0.06


Officials of 7 developed countries consider NP as pollution substances and started NO_x, NP elimination at around 1980. Then CO₂ assimilation is retarded. Food like grain, fish production is retarded. CO₂ fix is retarded. I am insisting NO_x NP elimination should be stopped many times (ref 1-36). In this paper, I wish to tell NO_x, NP elimination is giving very bad effect for the economy, global warming.(ref 19) .

Effect of NO_x, NP elimination on GWPR(global warming protection

ratio) and GDP

When 140 billion tone fossil is burned 420 billion tone CO₂ and 16.8 billion tone NO_x are produced. About 380 billion tone CO₂ is fixed by CO₂ assimilation. About 140 billion tone CO₂ must be reduced .Most of CO₂ can be reduced by CO₂ assimilation. We must promote CO₂ assimilation. We must provide enough NP fertilizer. NO_x and NP in waste water are best sources of NP fertilizer.

. Officials of developed countries put emphasis of toxicity than utility of NO_x, NP. They started elimination of NO_x by ammonia.



Amount of NO_x 16.8 billion tone is so much. 7 times of synthetic nitrogen fertilizer 2 billion tone of the world. To destroy one nitrogen fertilizer with one other nitrogen fertilizer is giving tremendous loss.

NO_x is very effective promotor of CO₂ assimilation. Therefore the production of grain and fish increased proportionally by the increase of CO₂ and NO_x. In 1900 20 billion tone CO₂ is emitted and 20 billion tone CO₂ is fixed. In 1960 100 billion tone CO₂ is emitted and 100 billion tone CO₂ is emitted and 100 billion tone CO₂ is fixed. In 1980 200 billion tone CO₂ is emitted and 180 billion tone CO₂ is fixed. In 2016 360 billion tone CO₂ is emitted and 220 billion tone CO₂ is fixed. Amount of CO₂ fix is 140 billion tone less than emission. This is caused by the elimination of NO_x and NP.

By the elimination of NO_x, CO₂ assimilation is retarded. Agriculture and fish industry of developed countries are declining.

CO₂em(CO₂ emission),NO_x(NO_x production),NO_xc(NO_x concentration at exit gas),GWPR(global warming protection ratio), GDP(GDP increase ratio) of 13 countries are shown in Table 1

Table1

Country	CO ₂ em bill t	NO _x bill t	NO _x con g/kWh	Area km ²	FixableCO ₂ bill t	GWPR inc ratio	GDP
World	360	14.4					
China	106.4	4.25	1.6	.1.0x 10 ⁷	100	1.0	6.9

USA	51.0	2	0.5	9.5x10 ⁶ 95	0.53	1.48
India	24.6	1	1.6	3.2x 10 ⁶ 32	0.76	7.1
Japan	12.5	0.5	0.1(2018) 1.6(1980)	3.8x 10 ⁵ 3.7	3.4 1.5	1.03 7.0
Russia	19.6	0.63		3.2x 10 ⁶ 32	0.61	0.8
Germany	7.8	0.31	1.0	3.5x 10 ⁶ 3.5	2.2	1.83
Iran	6.3	0.25		1.6x 10 ⁶ 1.6	3.9	2.6
Canada	5.6	0.22	1,3	1.0x 10 ⁸ 100	0.06	1.44
Indonesia	5.0	0.2	1.6	1.9x 10 ⁶ 19	0.3	5.2
U. K	4.0	0.16	1.3	2.4 x 10 ⁴ 2.4	1.7	1.8
Turkey	4.0	0.16		7.8x 10 ⁷ 7.8	0.5	-2
Italy	3.5	0.14	0.5	2.0x10 ⁵ 3.0	1.2	0.88
France	3.3	0.13		6.4x 10 ⁵ 8.4	0.4	1.2

! Km² green land can fix 1000 t CO₂. Fixable CO₂ of the country can be estimated by 1000 x area of the country.

Amount of NO_x produced at world is 16.8 billion tone. Developed countries are eliminating about 6 billion tone NO_x producing 10 billion tone CO₂. 6 billion tone NO_x can fix 6x 25 = 150 billion CO₂. Therefore if developed countries stop NO_x elimination, 150+ 10= 160 billion tone CO₂ emission is reduced and global warming can be protected.

When we look at high GWPR countries, Japan 3.4 Germany 2.2, Iran 3.9 , U.K 1.7, Italy 1.2 , These countries area are narrow and they cannot fix produced CO₂ at his countries.

Growth rate of GDP of the countries who eliminate NO_x are small as USA 1.46, Germany 1.83, Japan 1.03, Canada 1.44, U.K 1.6, Italy 0.88 .

At China 4.25 billion tone. USA 2 billion tone, India 1 billion tone, Japan 0.5 billion tone NO_x are produced. Japan eliminating this 0.5 billion tone. Butane 0.1280 billions is used for the production of H₂ 0.0606 billion tone and CO₂ 0.7480 billion tone is produced. If Japan stop NO_x elimination , 25 times of NO_x 0.5x 25= 12.5 billion tone CO₂ can be fixed.. By doing plankton CO₂ assimilation at 3 times area of Japan land, 3.8x 10⁵Km² area, 11.4 billion tone CO₂ can be fixed. 0.745 billion tone CO₂ by stopping of NO_x elimination can be saved. 0.5 billion tone CO₂ by stopping NP waste water purification can be saved. Total 11.4 + 0.745 + 0.5 = 12.645 billion tone CO₂ generation can be stopped. Japan can produce 0.3 billion fish and Japanese can enjoy anti-aging and long life. (Ref 26,37-42) If europa stop the elimination of 0.71 billion tone NO_x and 0.2 billion tone NP, 10 billion tone CO₂ generation can be stopped. And 0.1 billion tone fish can be produced

China producing 106.4 billion tone CO₂. Area of China is 1.0x 10⁷ km²

China can fix 100 billion tone CO₂. GWPR = 106.5/ 100 = 1.0

Low area country Japan GWRP = 12.5/ 3.7 = 3.4

NO_x elimination can be found by NO_x concentration of exit gas. 1.6 g/kwh is no elimination. 0.1 g/kWh is complete elimination No NO_x elimination countries like China, India, Indonesia show low GWPR and high GDP growth rate. On the contrary,

NO_x eliminating country like Japan (3.4 1.03), Germany (2.2 1.83) UK (1.7, 1.8), Italy (1.2 , 0.88) show high GRPR and low GDP growth rate.

Japan is eliminating NO_x , NP most severely. NO_x concentration at exit gas is 0.1 g/kWh. Then fish production decreased from 12 million tone in 1970 to 2 million tone In 1985 by NO_x NP elimination policy . And DGP do not increase for 40 years from 1980.

Low doses of inhalation of nitric oxide have been reported to be clinically effective, and most current dosing recommendation do not exceed 40 ppm. At this dose, the little measurable short term toxicity. Indeed, it is noteworthy that in the large randomized trials of inhalation of nitric oxide, major clinical toxicity (e.g. methemoglobinemia) was observed only at dose > 80 ppm (Ref 43,44). Therefore NO_x has small demerit but not significant as big merit that NO_x is essential for the growth of plant for the production of food for the promotion of health and long life. The ratio of merit / demerit is 10000/1. NO_x elimination at exit gas of factory and garbage incinerator should be stopped.

Japan producing too much CO₂

I investigated how much CO₂ is produced by the elimination of NO_x and NP in Japan

Japan emitting 9.1 tone CO₂ per person. This value is too many in compared with France 5.6 tone, UK 5.7, Italy 5.7. I found that Japan producing 2 billion tone CO₂ for the elimination of NO_x (Ref 29) and NP in drainage and elimination of NO_x at garbage incinerator exhaust gas (Ref 36).

NP elimination in waste water should be stopped

Japan constructed 2200 waste water purification stations to eliminate NP. Much CO₂ is produced for the construction of 2200 waste water purification stations.

I investigated Yamazaki waste water purification center at Yamazaki, Kamakura in Japan (ref 31). This center cover 96881 persons. Water 98287 m³ containing Nitrogen 40mg /l, Phosphorous 4.2mg/l is treated by activated sludge process. Air is bubbled for ten hours to give water containing Nitrogen 7.5 mg Phosphorous 2.7 31mg/l. Consuming 8841200 kWh electricity. This data showed that 7.34 Kg Nitrogen, 2.65 Kg Phosphorous is eliminated in one day at this center. This data indicate $7.34 \times 120000000 / 96881 \times 365 = 140$ million tone nitrogen, 12.8 million tone phosphorous are eliminated in Japan in one year. Population of Japan is 1.2 billion. $8841200 \times 120000000 / 96881 = 110$ billion kWh electricity is consumed in Japan for the treatment of waste water. This correspond $100880 / 110 = 1.11\%$ of total electricity consumption 100880 kWh of Japan

If waste water purification is not done in Japan, $1.40 \times 25 = 35$ million tone CO₂ is fixed and 35 million tone plankton can grow and $35 \times 1/10 = 3.5$ million tone fish will be produced.

Bon fire inhibition rule should be abandoned

In Japan waste material must burn at incinerator. 0.4289 billion tone garbage(331 kg per person) is produced. Japan constructed 1243 garbage incinerators. Top number in the world.

Second is USA 351 third France 181 . Japan reconstructed high temperature garbage incinerator in 2002. About 2 billion CO₂ is produced for construction of these garbage incinerator.

In Japan very special law about the garbage incinerator was set up in 2002 by the reason much NO_x is produced at lower temperature. By this rule, incinerator must be burned at higher temperature than 800 °C by adding excess fuel to keep higher temperature. Corrugated carton and fallen leaves must be burned at high temperature incinerator. Bon fire is inhibited by the reason bon fire produce much NO_x. Burning of rice straw wheat straw at rice field is not possible. Big earth quake and tsunami happened in east Japan in 2011. Debris disposal was not allowed to burn on site. Debris disposal must transfer to far away district having high temperature incinerator consuming much fuel and money. Operation of this high temperature incinerator is using much excess fuel releasing much CO₂. There is Nagoshi clean center at Kamakura ,Japan This clean center burn garbage 0.03 million tone at Kamakura producing 0.045 million tone CO₂. Exhaust gas contain NO_x. By insertion of ammonia This center used 40.94 kg ammonia in 2018. This mean $40.94 \times 30/17 = 72.256$ kg NO is eliminated by ammonia at Negoshi clean center. (ref 36). Population of Kamakura is 0.172 million. This data indicate $72.256 \times 120000000/172000 = 50.41$ million kg NO is eliminated at burning of garbage in Japan. $40.94 \times 12000/17.2 = 285.64$ million kg NO_x is eliminated by 255 million kg ammonia. 255 million kg ammonia is produced from 54 million kg H₂ . If NO_x elimination is not done 706 million kg CO₂ is not produced. 285 million kg NO_x can fix $0.285 \times 25 = 7.125$ million tone CO₂ .

The countries who use NO_x , NP are growing and increasing population. The countries who eliminate NO_x,NP are declining and decreasing population(Ref 32). DGP, food and population can be increased by effective use of NO_x and NP (ref 16,19,21,32-36)

Summary

Complete recycle of N and P is essential for complete recycle of CO₂

1. NO_x produced by burning should be released as it is. Do not eliminate NO_x with ammonia.

2. Close up waste water purification center. Excreta should be released as it is. Ocean dumping, river dumping, field dumping, agriculture field dumping, forest dumping are recommended.
3. Garbage should be burned on site. Kitchen waste should be buried .
4. Bon fire , slash and burn agriculture should be encouraged.
5. NOx elimination law should be abandoned
6. Waste water purification law should be abandoned
- 7 Bon fir inhibition law should be abandoned
8. Stop the unproductive spent of fossil fuel ,like war, military exercise .auto race, leisure cruising and leisure trip
- 9 Stop the unnecessary economy stimulus measure such as renewal of building, road
10. Restriction rule of NOx emission of car should be loosened

Reference

1. Ozaki Shoichiro Recycle of nitrogen and phosphorous for the increase of food production New Food Industry 1993 35, No 10 33-39.
- 2 .Ozaki Shoichiro Methods to protect global warming. Adv Tech Biol Med. 2016, 4. 181
3. Ozaki Shoichiro. Methods to protect global warming, Food production increase way. New Food Industry 2016 58 No 8 47-52
4. Ozaki Shoichiro Global warming can be protected by promotion of CO₂ assimilation using NOx. Journal of Climatology & Weather Forecasting 2016 4.2 1000171
5. Ozaki Shoichiro Global warming can be protected by promotion of plankton CO₂ assimilation. Journal of Marine Science: Research & Development 2016 6.213
6. Ozaki Shoichiro Method to protect global warming by promotion of CO₂ assimilation and method to reactivate fish industry. New Food Industry 2017 59 No 3 61-70
8. Ozaki Shoichiro Protection of global warming and burn out of fossil fuel by promotion of CO₂ assimilation . J.of Marine Biology & Oceanography 2017, 6:2 NOxU

9. Ozaki Shoichiro. Promotion of CO₂ assimilation supposed by NO_x is best way to protect global warming and food production. 2017, Artiv of Pet-EnvironBiotechnol 02.110
10. Ozaki Shoichiro. Promotion of CO₂ assimilation supported by NO_x is best way to protect global warming. J.Marine Biol Aquacult. 2017 vol 3. Issue 2.
11. Ozaki Shoichiro Stopping of NO_x elimination is easy way to reduce CO₂ and protect global warming J.Environ Sci Public Health 2017:1 (1)]24-34
12. Ozaki Shoichiro Stopping of NO_x elimination is clever way to reduce CO₂ and to increase fish production J.of Cell Biology 6 Immunogy 2017 1 e 102
13. Ozaki Shoichiro Effective uses of NO_x and drainage are clever way to protect global warming and to increase fish production. Oceanography & Fisheries.2017 4(4)
14. Ozaki Shoichiro NO_x Elimination and Drainage NP Elimination should be stopped for the production of fish and for the protection of global warming . J.of Fisheries and Aquaculture Development 2017 issue 05 125
15. Ozaki Shoichiro Let`s enjoy civilized life using limited amount of fossil fuel. Journal of Aquaculture & Marine Biology 2017 6 (3) 06 00158
16. Ozaki Shoichiro Method to fit Paris agreement for protection of global warming . International Journal of Waste Resources 2017 7-4 318 doi:10.4172/2252-5211.1000318
17. Ozaki Shoichiro Method to protect global warming and to produce much fish by promotion of plankton growth New Food Industry 2018 60 no3 88-94
18. Ozaki Shoichiro Method to protect global warming by promotion of plankton CO₂ assimilation Rikuryou Science .2018 61 23
19. Ozaki Shoichiro Effect of NO_x elimination on electricity price, fish production,GDP and protection of global warming International J of Waste Resources 2018 8 issue 1 1000328 doi:10.4172/2252-1000328
20. Ozaki Shoichiro How to fix carbon dioxide same amount as emission for the protection of global warming Research & Development in Material Science 2018 vol 3 issue 5 RDMS.000572
21. Ozaki Shoichiro Stop of NO_x elimination and stop of wast water purification are easy methods to protect global warming 2018 J of Immunology and Information Diseases Therapy 1 1 [doi.org/06.2018/1.10006](https://doi.org/10.4172/2252-1000328)
22. Ozaki Shoichiro Climate can be regulated by effective use of NO_x and wast water NP 2018 Biomedical Research and Reviews volume 1.1
23. Ozaki Shoichiro Promotion of Plankton CO₂ assimilation by effective use of NO_x and NP is best method to produce much fish and protect global warming 2018 J of Marine Science Research and Oceanography Volume 1 issue 1. 1
24. Ozaki Shoichirou Promotion of plant growth by NO_x is best method to reduce CO₂ and to protect global warming and to get best climate. International J. of Earth and environmental Science 2018 3 160

- .25. Ozaki Shoichiro Promotion of plant growth by NO_x is best method to reduce CO₂ and to protect global warming Current Trends in Oceanography and Marine Science 2018 01 1-4
- 26 Ozaki Shoichiro Fish is best food to get anti-aging and long life. NO_x elimination should be stopped to produce much fish and to protect global warming Jacobs J of physiology 2018 4(1) 017
- 27 Ozaki Shoichiro Fish is Best Food to Get Anti-Aging and Long Life J of Aging and Neuropsychology 2018 issue 2 1-6 DOI: <http://dx.doi.org/10.20431/2454-7670.0501001>
- 28 Ozaki Shoichiro NO_x and NP in waste water fix CO₂ and control global warming and climate International J of Biochemistry and Physiology 2018 3 (4) DOI: 10.23880/ijbp-16000140
29. Ozaki Shoichiro Why global warming is progressing. Promotion of CO₂ assimilation is best method to protect global warming Rikuryou Science .2019 62 16-18
- 30 Ozaki Shoichiro The effect of of increase of NO_x and CO₂ on grain and fish production , protection of global warming and climate International Journal of Earth Science and Geology 2019 1(1) 6-10
31. Ozaki Shoichiro Complete use of NO_x and NP is essential for the increased production of food and protection of global warming. Inter.J. Innovative Studies in Aquatic Biology and Fisheries 2019 3 (1) 1-6
32. Ozaki Shoichiro Increase of CO₂ and NO_x promote CO₂ assimilation, CO₂ fix and food production Advances in Bioengineering & Biomedical Science Research 2019 2 issue 3 1-6
33. Ozaki Shoichiro: Promotion of CO₂ assimilation by effective use of NO_x and NP is best method to produce much fish and protect glow warming *EC Agriculture* 2019 5: Issue 8, 492-497.
- 34 Ozaki Shoicjiro. Why fish production of Japan decreased. Why global warming is progressing. New food Industry 2019 Vol 61 No 10 787-793
- 35 Ozaki Shoichiro In pure water no fish can live. Water purification promote global warming, decline of countries. Rikuryou Science .2020 63
36. Ozaki Shoichiro NO_x elimination and NP elimination are promoting global warming *EC Agriculture* 2019
37. Ozaki Shoichiro. Synthesis of anti-ageing reagent: Sulfo disaccharide co-working with anti-aging gene. *Archeves of Medicines* (2015) 7.No.6:17
38. Ozaki Shoichiro . Sulfo disaccharides co-working with Klotho. Studies on structure , structure activity relation and function. *World J of Pharmacy and Pharmaceutical Sciences* (2015) 4 152-175

39. Ozaki Shoichiro . Nutrition for good health, anti-aging and long life. Hyaluronic acid,glucosamine and chodroitin. Maternaland Paediatric Nutrition Journal (2015) 1: e102.
- 40.Ozaki Shoichiro Food containing hyaluronic acid and chondroitin is essential for anti-aging International J of Aging & Clinical Research 2016 1.101 <http://dx>.
41. Ozaki Shoichiro , Toward anti-aging and long life. Jacobs Journal of Physiology. (2016) 2(1); 12
- 42.Ozaki Shoichiro Secret of anti-aging: Anti-aging food containing glucosamine, hyaluronic acid and chondroitin. Jacobs Journal of Physiology (2016) 2(1): 13-17
- 43.Davidson D et al, Inhaled nitric oxide for the early treatment of persistent pulmonary hypertension . The 1-NO/PPHN Study Group. Pediatrics 1998 101, 325-334.
44. Barry Weinberger, Debra L. Laskin, Diane IE.Heck and Jeffrey D.Laskin, The Toxicity of inhaled Nitric oxide. Toxicology Science 2000 59,1,3-16.