

## Microsculpture of Nutlets Surface of some Libyan *Salvia* L. species (Lamiaceae)

Ghalia T. El Rabiai and Khatria K. Elfaidy

Department of Botany, Faculty of Science, Benghazi University, Libya

### ABSTRACT

A comprehensive morphological and micro-morphological study of the nutlets of five Libyan *Salvia* species (Lamiaceae) was conducted to evaluate nutlets characteristics by using scanning electron microscopy (SEM). Differences in surface ornamentation, size, shape and color were observed between the species. The studied species were categorized in three basic types based on surface ornamentation: irregular prominences, regular prominences and smooth nutlets. The shape of nutlets were described as oblong, ovoid-oblong to globose-subglobose and their size range is 2-3.5 mm in length and 1.5-2.5 mm in width. Nutlet micromorphological characteristics such as surface ornamentation can be useful for classification and identification of *Salvia* species in Libya.

**Keywords:** *Salvia*, Microsculpture, Nutlets, Lamiaceae, Libya

### INTRODUCTION

The genus *Salvia* L. (Lamiaceae) is one of the largest genera in this family (Cvetkovikj et al. 2015). The plant name *Salvia* (sage) comes from the Latin word *salvare*, which means healer (TOPÇU et

al. 2013). The genus *Salvia* L. belongs to the Mentheae tribe within the Nepetoideae subfamily (Kharazian 2014) includes around 1000 species that have almost cosmopolitan distribution (Saravia et al. 2018); In Libya, it is represented by 10 species; out of which 3 are cultivated (Jafri, 1985). Numerous species of the *Salvia* genus are economically important since they are used as spices and flavouring agents in the field of perfumery and cosmetics (Felice Senatore et al., 2004 and 2006); and some species of *Salvia* have been cultivated worldwide for use in folk medicines (Tohamy et al. 2012). Nutlet ornamentation, shape, size and colour particularly proved to be good taxonomic characters for the Egyptian Lamiaceae taxa (Kamel, 2014).

*Salvia* L. has been shown that gross morphology of nutlets and their sculpturing pattern are variable and taxonomically useful at a species level (Oran 1996). Studies on nutlet micromorphology within Lamiaceae showed that nutlets features e.g., shape and surface sculpturing, were potentially useful at different taxonomic levels (Moon et al., 2009; Khosroshahi & Salmaki, 2018). Among different nutlet characters, type of sculpturing has been considered to be taxonomically most important (Kahraman et al., 2011), however, color, size and shape of nutlets were considered unimportant, either because they did not vary or the variation was random or too great (Oran, 1996). Ozkan et al., reported the nutlets are placed in three groups based on the shape and ornamentation. Ozkan explained *S. aethiopsis* and *S. virgata* nutlets ornamentation are foveate and

reticulate, While Mousavi, S. et al (2013) preferred to name, surface with hexagonal prominences and undulated stripped respectively.

Recent studies of the mericarp morphology of Lamiaceae taxa have contributed useful information at different taxonomic levels. The importance of mericarp morphology in the modern taxonomy of Lamiaceae has been emphasized by many researchers (Kaya & Dirmenci 2012, Dinç et al. 2009 and Kaya et al. 2014).

Kahraman & Dogan (2010) reported that the pollen size, shape and exine ornamentation, and nutlet micromorphology in the genus *Salvia* were important in distinguishing between the species.

## **MATERIAL AND METHODS**

Specimens from five *Salvia* species were collected from several localities in Libya, collected from different localities of Libya between January 2019 and October 2020 . Identification of species was performed using Flora of Libya (JAFRI et a 1985), Flora Of Europaea (Tutin et al 1992). Nutlets (Mericarps) were first observed with a stereomicroscope to ensure that they were of normal size and maturity. For nutlets length and width, five samples of each taxon were taken and measured. For Scanning Electron Microscope (SEM) observation, dried mature nutlets of five individuals for each species were examined using a SEM microscope model (Joel, JFC 1100). Nutlets were directly mounted on aluminum stubs using

double-sided adhesive and were sputter coated with a thin layer of gold. Coated nutlets were examined and SEM micrographs were captured at 200× and 500×. The terms used for describing the cypsela surface patterns have been adopted according to Stearn (1992). All photographs were taken at Central Laboratory of Alexandria University, Alexandria, Egypt. (Figure 1).

## Results










In this study, macromorphological and micromorphological characters of Nutlets were observed via SEM. Nutlets shape, size and other features of Nutlets were given in table (1). In addition, illustrating SEM photomicrographs are presented in Figure. (1). The shape was variable among the studied species; it is globose-subglobose, ovoid and oblong. The colour of nutlet was Pale green with netted lines brown, dark brown and black. The mean length of mature nutlets of the studied species ranged from 2 mm to 3.5 mm. The smallest nutlet recorded in *S. verbenaca* whereas nutlet of *S. fruticosa* and *S. viridis* where the longest ones. The observations in the studied species, surface patterns were divided in three types: **Type I:** Regular: regular hexagonal prominences as in *S. fruticosa* and regular prominence in chain form as in *S. viridis*. **Type II:** Irregular prominence with fine and rough folds as in *S. lanigira* and Irregular prominence with deep and shallow channels as in *S. verbenaca*.

**Type III:** No microphological features observed (smooth) as in *S. spinosa*.


**Table 1.** Macro- and micromorphological features of the studied species

Species	Shape	Colour	Length (mm)	Width (mm)	Ornamentation
<i>S. fruticosa</i>	Globose-subglobose	Dark brown to black	3 - 3.5	2.5	Regular reticulate, hexagonal, with narrow depressed anticlinal wall and wide raised periclinal wall
<i>S. lanigira</i>	Ovoid, oblong	Dark brown to black	2.25-2.5	1.25-1.5	Irregular prominence with fine and rough folds
<i>S. spinosa</i>	Ovoid	Pale green with netted lines	3	2	No microphological features observed (smooth)
<i>S. verbenaca</i>	Ovoid	Dark brown	2	1.5	Irregular prominence with deep and shallow channels
<i>S. viridis</i>	Oblong	Brown	3-3.5	2	Regular prominences in chain form with tangled

					strands on the surface
--	--	--	--	--	------------------------



**Fig.(1):** Scanning electron micrographs of Nutlets in *Salvia* species consist of 1-shape (x25) ; 2 surface ornamentation (x500): a- *S. fruticosa*; b- *S. lanigera* ; c- *S. spinosa*.



**Fig. (1):** Scanning electron micrographs of Nutlets in *Salvia* species consist of 1-shape (x25); 2 surface ornamentation (x500): D- *S. verbenaca*, E- *S. vrbenaca*.

## Discussion

Ozkan et al. (2009) reported that the nutlets were placed in three groups based on the shape and ornamentation (spherical, trigonous and prolate spheroidal) and (foveate, reticulate and verrucate)

respectively. Between 12 studied *Salvia* nutlets in their study, *S. ceratophylla*, *S. aethiopsis* and *S. virgata* were common with the present research. Ozkan explained *S. aethiopsis* and *S. virgata* nutlets ornamentation are foveate and reticulate while in present research, it was preferred to name, surface with hexagonal prominences and undulated stripped respectively. Also, Kahraman et al.(2009) pointed the size, shape and ornamentation of *S. ballsiana*, *S. macrochlamys* and *S. hedgeanaare* diagnostic. Our findings agree with those of previous studies in Lamiaceae confirming the usefulness of Nutlet characteristics. Nutlet morphology provides valuable data for delimitation of closely related species

In conclusion, the shape, color and ornamentation of *Salvia* nutlets varied among the species and those are taxonomical characters help to identify species.

## References:

- Cvetkovikj, I., Stefkov, G., Karapandzova, M., & Kulevanova, S. (2015). Essential oil composition of *Salvia fruticosa* Mill. populations from Balkan Peninsula. Macedonian pharmaceutical bulletin, 61(1), 19-26.
- Dinç, M., Dođiu, S., Bilgili, B. & Duran, A. (2009) Comparative anatomical and micromorphological studies on *Teucrium creticum* and *Teucrium orientale* var. *orientale* (*Teucrium* section *Teucrium*, Lamiaceae). nordic Journal of Botany 27: 251-256.



- Felice S., N. A. Arnold, F. Piozzi and C. Formisano. (2006). Chemical composition of the essential oil of *Salvia microstegia* Boiss. et Balansa growing wild in Lebanon. J Chromatogr A.; 1108: 276-278.
- Felice S., N. A. Arnold and F. Piozz. (2004). Chemical composition of the essential oil of *Salvia multicaulis* Vahl. var. *simplicifolia* Boiss. J Chromatogr A., 1052: 237-240.
- JAFRI, S.M.H & EI-GADI (1985). A. Flora of Libya, (Lamiaceae), Al-faateh University , Faculty of science , Department of Botany , Tripoli-Libya , Vol.118 ,.
- Kahraman, A., Celep, F., & Dogan, M. (2009). Comparative morphology, anatomy and palynology of two *Salvia* L. species (Lamiaceae) and their taxonomic implications. Bangladesh Journal of Plant Taxonomy, 16(1), 73-82.
- Kahraman A, Celep F, Dođan M, Guerin GR, Bagherpour S. (2011). Mericarp morphology and its systematic implications for the genus *Salvia* L. section *Hymenosphace* Benth.(Lamiaceae) in Turkey. Plant systematics and evolution.;292:33-9.
- Kamel, W. (2014). Nutlet morphology and its taxonomic implication in some taxa of Lamiaceae in Egypt. *Taeckholmia*, 34(1), 101-127.
- Kaya, A. & Dirmenci, T. (2012) Nutlet morphology of Turkish *Ziziphora* L. (Lamiaceae). *Plant Biosystems* 146: 560-563.
- Kaya, A., Dirmenci, T. & Satıđl, F. (2014) Morphological studies on the nutlet of Turkish *Cyclotrichium* Manden. & Scheng. (Lamiaceae). *Plant Biosystems* 149: 984-989.
- Kharazian, N. (2014). Chemotaxonomy and flavonoid diversity of *Salvia* L.(Lamiaceae) in Iran. *Acta Botanica Brasilica*, 28(2), 281-292.

- Khosroshahi, E. and Y. Salmaki (2018). Nutlet micromorphology and its systematic implications in *Phlomis* Moench(Lamiaceae), *Nova Biologica Reperta* 5(1):82-94
- Moon H-K, Hong S-P, Smets E, Huysmans S. (2009.) Micromorphology and character evolution of nutlets in tribe Mentheae (Nepetoideae, Lamiaceae). *Systematic botany*.34:760-76.
- Mousavi S. M., A. Jafri and S. Najafi (2013). Nutlet Micromorphological Study on *Salvia* L. (Lamiaceae) from NE Iran. *American Journal of Plant Sciences*, 4, 1457-1460.
- Oran, S. A., (1996). Ultrastructure of nutlet surface of the genus *Salvia* L. in Jordan and the neighbouring countries. *Dirasat, Natural and Engineering Sciences* 23, 393-408.
- Özkan, M., Aktaş, K., Özdemir, C., & Guerin, G. (2009). Nutlet morphology and its taxonomic utility in *Salvia* (Lamiaceae: Mentheae) from Turkey. *Acta Botanica Croatica*, 68(1.), 105-115.
- Saravia, A., & Pinto, C. (2018). Pollen morphology of four species of *salvia* genus (lamiaceae) in periurban areas of Sucre, Bolivia. *Revista Ciencia, Tecnología e Innovación*, 16(17), 1013-1017.
- Stearn W.T. (1992). *Botanical Latin*. 4<sup>th</sup> edition. David & Charles Publishers, London, pp. 489-491.
- Tohamy, A. A., Ibrahim, S. R., & Moneim, A. E. A. (2012). Studies on the effect of *Salvia aegyptiaca* and *Trigonella foenum graecum* extracts on adult male mice. *Journal of Applied Pharmaceutical Science*, 2(5), 36.
- Topcu, G., Öztürk, M., Kusman, T., Demirköz, A. A. B., Kolak, U., & Ulubelen, A. (2013). Terpenoids, essential oil composition, fatty acid profile, and biological activities of Anatolian *Salvia fruticosa* Mill. *Turkish Journal of Chemistry*, 37(4), 619-632.



Tutin T.G., V. H. Hewood, N.A. Burges, D.M. Moore, D.H. Valentine, S.M. Walters and D. A. Webb (1992). *Flora Urobaea. Lamiaceae* Vol. 3, Cambridge University Press.